



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

where g is the force of gravity, and σ , which is supposed to be greater than ρ , the density and a the radius of the sphere.

Hence,

$$6\pi\mu'\rho aV = \frac{4}{3}\pi g a^3(\sigma - \rho),$$

or

$$V = \frac{2}{9} \frac{g}{\mu'} \left(\frac{\sigma}{\rho} - 1 \right) a^2 = \frac{2}{9} \frac{g a^2 \sigma}{\eta},$$

by placing

$$\frac{\sigma - \rho}{\mu\rho} = \frac{\sigma}{\eta}.$$

QUESTIONS AND DISCUSSIONS.

SEND ALL COMMUNICATIONS TO U. G. MITCHELL, University of Kansas.

DISCUSSIONS.

CONCERNING TWO FIFTH-POWER PROBLEMS IN DIOPHANTINE ANALYSIS.

By CYRUS B. HALDEMAN, Ross, Butler County, Ohio.

I. To resolve

$$p^5 + q^5 + p^5 + s^5 = t^5 + u^5 + v^5 \quad (1)$$

where p, q, r, s, t, u , and v are rational.

Solution. Let $p = 2x$, $q = y - x$, $r = 2a - b^2 - x$, $s = 2a + b^2 + x$, $t = x + y$, $u = 2a - b^2 + x$ and $v = 2a + b^2 - x$.

Substituting in (1) we get

$$\begin{aligned} (2x)^5 + (y - x)^5 + (2a - b^2 - x)^5 + (2a + b^2 + x)^5 \\ = (x + y)^5 + (2a - b^2 + x)^5 + (2a + b^2 - x)^5 \end{aligned} \quad (2)$$

We may write (2):

$$\begin{aligned} (x + y)^5 + (x - y)^5 + [(2a + b^2) - x]^5 - [(2a + b^2) + x]^5 \\ + [(2a - b^2) + x]^5 - [(2a - b^2) - x]^5 - (2x)^5 = 0. \end{aligned} \quad (3)$$

Expanding, regarding each binomial in parenthesis as a single quantity, and adding, we have from (3)

$$\begin{aligned} 2x^5 + 20x^3y^2 + 10xy^4 - 2x^5 - 20x^3(2a + b^2)^2 - 10x(2a + b^2)^4 \\ + 2x^5 + 20x^3(2a - b^2)^2 + 10x(2a - b^2)^4 - 32x^5 = 0. \end{aligned} \quad (4)$$

Equation (4) reduces to

$$2x^2y^2 + y^4 - 16ab^2x^2 - 64a^3b^2 - 16ab^6 - 3x^4 = 0.$$

Solving for y^2 , we get

$$y^2 + x^2 = \sqrt{4x^4 + 16ab^2x^2 + 64a^3b^2 + 16ab^6} = 2x^2 + 4a(b^2 + d^2), \text{ say.}$$

From this we find

$$x^2 = \frac{4a^2b^2 - (b^2 + d^2)^2a + b^6}{d^2},$$

and

$$y^2 = \frac{4a^2b^2 - (b^2 + d^2)^2a + b^6 + 4d^2(b^2 + d^2)a}{d^2}.$$

To rationalize the expressions for x and y , let

$$d^2x^2 = 4a^2b^2 - (b^2 + d^2)^2a + b^6 = (n - 2ab)^2, \quad (5)$$

and

$$d^2y^2 = 4a^2b^2 - (b^2 + d^2)^2a + b^6 + 4d^2(b^2 + d^2)a = (n - 2ab - Z)^2. \quad (6)$$

From (5) we get

$$a = \frac{n^2 - b^6}{4bn - (b^2 + d^2)^2},$$

and from (6), since the square of $n - 2ab$ is assumed equal to the first four terms of the left-hand member, we have

$$a = \frac{2nZ - Z^2}{4bZ - 4d^2(b^2 + d^2)}.$$

Equating the values of a and clearing of denominators, we get

$$\begin{aligned} 4b^6d^2(b^2 + d^2) - 4b^7Z - 4d^2(b^2 + d^2)n^2 \\ = Z^2(b^2 + d^2)^2 - 4bZ^2n + 4bZn^2 - 2Z(b^2 + d^2)^2n. \end{aligned}$$

Place $4bZn^2 = -4d^2(b^2 + d^2)n^2$ and solve for

$$Z = -\frac{d^2(b^2 + d^2)}{b};$$

then

$$n = \frac{4b^7Z - 4b^6d^2(b^2 + d^2) + Z^2(b^2 + d^2)^2}{4bZ^2 + 2Z(b^2 + d^2)^2}.$$

Substituting the value of Z in n gives

$$n = \frac{d^2(b^2 + d^2)^3 - 8b^8}{2b(d^4 - b^4)}.$$

The value of Z and the value of n in terms of b and d , substituted in either of the above values of a gives

$$a = \frac{d^4(b^2 + d^2)^2 - 4b^8}{4b^2(d^4 - b^4)}.$$

From (5) and (6) we have

$$x = \frac{n - 2ab}{d},$$

and

$$y = \frac{n - 2ab - Z}{d}.$$

The values of a , n and Z substituted in these expressions for x , and y , give

$$x = \frac{bd^2(b^2 + d^2)^2 - 4b^7}{2d(d^4 - b^4)} \quad \text{and} \quad y = \frac{bd^2(b^2 + d^2)^2 - 4b^7}{2d(d^4 - b^4)} + \frac{d(b^2 + d^2)}{b}.$$

From the values of a , x and y we obtain

$$\begin{aligned} p &= \frac{bd^2(b^2 + d^2)^2 - 4b^7}{d(d^4 - b^4)}, & q &= \frac{d(b^2 + d^2)}{b}, \\ r &= \frac{2b^4d(b^4 - d^4) + d^2(b^2 + d^2)^2(d^3 - b^3) + 4b^8(b - d)}{2b^2d(d^4 - b^4)}, \\ s &= \frac{2b^4d(d^4 - b^4) + d^2(b^2 + d^2)^2(b^3 + d^3) - 4b^8(b + d)}{2b^2d(d^4 - b^4)}, \\ t &= \frac{d^4(b^2 + d^2) - 4b^8}{bd(d^4 - b^4)}, \\ u &= \frac{2b^4d(b^4 - d^4) + d^2(b^2 + d^2)^2(b^3 + d^3) - 4b^8(b + d)}{2b^2d(d^4 - b^4)}, \\ v &= \frac{2b^4d(d^4 - b^4) + d^2(b^2 + d^2)^2(d^3 - b^3) + 4b^8(b - d)}{2b^2d(d^4 - b^4)}. \end{aligned}$$

By substitution of the expressions just found for p , q , r , s , t , u and v in (1); and after clearing of denominators, we have the identity

$$\begin{aligned} &[2b^3d^2(b^2 + d^2)^2 - 8b^9]^5 + [2bd^2(b^2 + d^2)(d^4 - b^4)]^5 \\ &\quad + [2b^4d(b^4 - d^4) + d^2(b^2 + d^2)^2(d^3 - b^3) + 4b^8(b - d)]^5 \\ &\quad + [2b^4d(d^4 - b^4) + d^2(b^2 + d^2)^2(b^3 + d^3) - 4b^8(b + d)]^5 \\ &= [2bd^4(b^2 + d^2)^2 - 8b^9]^5 \\ &\quad + [2b^4d(b^4 - d^4) + d^2(b^2 + d^2)^2(b^3 + d^3) - 4b^8(b + d)]^5 \\ &\quad + [2b^4d(d^4 - b^4) + d^2(b^2 + d^2)^2(d^3 - b^3) + 4b^8(b - d)]^5 \end{aligned}$$

Take $b = 1$, $d = 2$, and we have, after dividing by 12^5 ,

$$16^5 + 50^5 + 53^5 + 79^5 = 63^5 + 66^5 + 69^5.$$

II. To resolve the equality

$$\begin{aligned} p^5 + q^5 + r^5 + s^5 + t^5 + u^5 + v^5 + w^5 \\ = (2g_1)^5 + (2g_2)^5 + (2g_3)^5 + \dots + (2g_n)^5, \end{aligned} \quad (1)$$

where $p, q, r, s, t, u, v, w, g_1, g_2, g_3, \dots, g_n$ are rational.

Solution. Let $p = a + b + c$, $q = a - b - c$, $r = b - a - c$, $s = c - a - b$,
 $t = d + e + f$, $u = d - e - f$, $v = e - d - f$, $w = f - d - e$.

Then, by substitution in (1), we have

$$\begin{aligned} & (a + b + c)^5 + (a - b - c)^5 + (b - a - c)^5 + (c - a - b)^5 \\ & \quad + (d + e + f)^5 + (d - e - f)^5 + (e - d - f)^5 + (f - d - e)^5 \quad (2) \\ & = (2g_1)^5 + (2g_2)^5 + (2g_3)^5 + \dots + (2g_n)^5. \end{aligned}$$

For convenience, we may write (2):

$$\begin{aligned} & [a + (b + c)]^5 + [a - (b + c)]^5 + [(b - c) - a]^5 - [(b - c) + a]^5 \\ & \quad + [d + (e + f)]^5 + [d - (e + f)]^5 + [(e - f) - d]^5 - [(e - f) + d]^5 \quad (3) \\ & = (2g_1)^5 + (2g_2)^5 + (2g_3)^5 + \dots + (2g_n)^5. \end{aligned}$$

Expanding, regarding each binomial in parenthesis as a single quantity and adding, we get from (3)

$$\begin{aligned} & 2a^5 + 20a^3(b + c)^2 + 10a(b + c)^4 - 2a^5 - 20a^3(b - c)^2 - 10a(b - c)^4 \\ & \quad + 2d^5 + 20d^3(e + f)^2 + 10d(e + f)^4 - 2d^5 - 20d^3(e - f)^2 - 10d(e - f)^4 \quad (4) \\ & = 32(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5). \end{aligned}$$

Equation (4) reduces to

$$\begin{aligned} & 10a^3[(b + c)^2 - (b - c)^2] + 5a[(b + c)^4 - (b - c)^4] \\ & \quad + 10d^3[(e + f)^2 - (e - f)^2] + 5d[(e + f)^4 - (e - f)^4] \\ & = 16(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5); \end{aligned}$$

and by further expansion and reduction, we have

$$5(a^3bc + ab^3c + abc^3 + d^3ef + de^3f + def^3) = 2(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5). \quad (5)$$

Let $d = a$ and (5) becomes

$$5a^3(bc + k^3ef) + 5a(b^3c + bc^3 + ke^3f + kef^3) = 2(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5).$$

Place $bc + k^3ef = 0$ and have

$$5a(b^3c + bc^3 + ke^3f + kef^3) = 2(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5).$$

From these equations we obtain

$$b = -\frac{k^3ef}{c}, \quad a = \frac{2(g_1^5 + g_2^5 + g_3^5 + \dots + g_n^5)}{5(b^3c + bc^3 + ke^3f + kef^3)},$$

which give the requirements.